Executive Summary: Air Quality in Our Area

Air Quality in Crawley

Air pollution is associated with a number of adverse health impacts. It is recognised as a contributing factor in the onset of heart disease and cancer. Additionally, air pollution particularly affects the most vulnerable in society: children, the elderly, and those with existing heart and lung conditions. There is also often a strong correlation with equalities issues because areas with poor air quality are also often less affluent areas[1,2].

The mortality burden of air pollution within the UK is equivalent to 28,000 to 36,000 deaths at typical ages[3], with a total estimated healthcare cost to the NHS and social care of £157 million in 2017[4].

As part of the Local Air Quality Management process (LAQM) required by the Environment Act 1995, the Council carries out an annual review and assessment of air quality in the borough to identify local air quality hot spots and pollution sources. Road traffic is the main source of (nitrogen dioxide) pollution in Crawley, and our network of monitoring sites measures concentrations along busy roads as well as at background locations and areas of specific interest (such as residential locations close to the airport), to give a broad picture of pollution levels across the borough.

Air Quality in Crawley is mainly good, with the exception of a small number of locations alongside busy roads where an air quality management area (AQMA) has been declared. The Council is working with its partners to produce an action plan (AQAP) which sets out ways to tackle pollution and target measures to improve air quality in these areas.

There are two national objectives for nitrogen dioxide (NO_2). These are for the average level over a year, which should be below 40 g/m^3, and the average level for one hour,

---

[1] Public Health England. Air Quality: A Briefing for Directors of Public Health, 2017

[2] Defra. Air quality and social deprivation in the UK: an environmental inequalities analysis, 2006

[3] Defra. Air quality appraisal: damage cost guidance, July 2021

[4] Publi

which should be below 200 g/m^3. Most of our NO_2 monitoring is done using diffusion tubes which can't measure hourly average, so guidance developed by the Department for Environment, Food and Rural Affairs (DEFRA) estimates that if the yearly average is below 60 g/m^3 it will meet the hourly objective of 200 g/m^3. The annual average objective applies to residential, hospital and educational sites. The hourly average objective applies to these sites as well as busy streets and workplaces.

The measured results for 2021 NO_2 are set out in tables A.4 and B.1 of this report and show that national air quality objectives were met at all of Crawley's monitoring sites (although one site in the AQMA near Three Bridges station was borderline). The 2021 levels of nitrogen dioxide were generally slightly higher than in 2020, when Covid travel

with Sussex-air and Sustrans the project completed its third phase in 2021/22, engaging with 55,000 young people at 83 schools across the County. Crawley primary schools made up 45% of the new schools joining the project in 2021.

No monitoring sites within the monitoring network exceeded the air quality annual and one-hour mean objectives for NO_2 in 2021.

There were no exceedances of the annual mean and 24-hour objectives for PM_{10} in 2021 and annual mean $PM_{2.5}$ was well below the national limit value of 25ugm^3. The 5-year trend for both PM_{10} and $PM_{2.5}$ continues to be down.

Although there were no exceedeances of the AQO for NO

car journeys. A large percentage of local traffic are journeys of less than 2km, and 58% of car trips are under 5km. These high volumes of traffic on our local roads cause congestion and contribute to worsening air quality. However, since many journeys are short, there is opportunity to improve local air quality by switching to sustainable transport options such as walking, cycling, public transport or car sharing.

Many of our action plan measures include schemes that are aimed at infrastructure improvements to reduce congestion, improve sustainable transport options and encourage modal shift. Awareness raising to support behavioural change is an important part of this process, and local engagement is integral to, not only understanding the needs of the community, but also for informing the public and stakeholders of proposed improvements.
In developing our air quality action plan it is important to work closely with interested parties, including Community groups, Elected Members, Transport Planners, Planning Policy and Development control. Although Covid restrictions have prevented some of

2.2 Progress and Impact of Measures to address Air Quality in Crawley

5. Council has plans to renew and update the AQAP based on the extension of the AQMA. It is understood that COVID-19 has caused hindrance due to limited resources within the team. Any progress with the AQAP should be reported in next year's ASR.

***Response**: The Council is actively engaged in tackling air quality issues and promoting sustainable strategies to improve air quality within the borough. Section 2.2 outlines in detail the measure we have progressed over the reporting year. However, progress on the action plan is still behind schedule.*

6. A detailed COVID report has been submitted alongside the report. This report breaks down the trends in pollutant levels in different areas within the Council and shows reduction in traffic volumes by approximately 60% in the region. The report supports maintaining the lockdown behavioural change in a sustainable way by means of reducing vehicle use and adopting alternative forms of transport such as cycling and walking. The Council is already working on some of these measures which is commended.

***Response**: The Council is continuing its work through the Crawley Growth Programme and LCWIP plans to improve sustainable transport infrastructure and encourage active travel.*

7. Hazelwick AQMA's date of amendment has not been updated on DEFRA's website. Please therefore update as appropriate.
   The AQMA name on Table 2.1 needs to be updated to 'Hazelwick AQMA' instead of Crawley AQMA.

Response:

reflect the wider area of than just the locality of Hazelwick.

The

submitted to Defra. At the time of writing this had not been updated on the Defra AQMA website, but email response received from LAQM-portal admin team has confirmed that the amendment will be published within the next update cycle of their website.

8. The AQMA map presented in Figure D.1 needs to be updated to show the

WSCC Electric Vehicle Strategy 2019-2030. The strategy aims to encourage and support the adoption of electric vehicles within the county. The objectives are to ensure sufficient charging infrastructure in place to support the vehicle charging and ensure renewable energy sources are used.

Since 30% of households in Crawley have no access to off road parking this creates a barrier to switching to EV. As a result, Crawley is identified in the strategy as a priority area for EV infrastructure.

https://www.westsussex.gov.uk/media/13766/electric_vehicle_strategy.pdf

During the reporting year 2021/22 the Council, in partnership with WSCC, commissioned Connected-Kerb to deliver the EV infrastructure network across the

The Project was delivered on behalf of the Sussex authorities by Sustrans, working with schools across Sussex to raise awareness of air pollution, and examine local solutions such as reducing traffic and replacing short journeys with active travel on “clean air routes” to avoid exposure to air pollution.

To date the project has engaged with 55,000 young people at 83 schools across the County. In 2021/22, 45% of primary schools joining the project were from Crawley.

Key outcomes from measure: Awareness raising/ behavioural change/modal shift

Crawley Climate Emergency Action Plan, ‘Action to Zero’, published on 24 November 2021 following Cabinet approval. The CEAP sets out how we will meet our carbon reduction targets.

Key outcomes from measure: Target to reduce emissions by 50%, and as close to net zero as possible by 2030 and to reach net zero by 2040. Reduced emissions also benefit background levels of NO_2

CBC/Net Zero Collective Group Research Project – working with Southampton university to survey and monitor the retrofit of 10 Crawley council properties (from mixed housing stock) to identify the most efficient and effective methodology for the retro fit and monitor the level of energy savings/carbon reduction in residential accommodation. The pilot project retrofit was completed 2022, and although still at monitoring stage, early findings have identified energy savings of at least 25%. And refined a retrofit methodology that conforms to Pas 2035 standards. The pilot will continue for a further 25 properties in 2022.

Key outcomes from measure: Reduce emissions, carbon reduction and NO_2 resulting in improved air quality

Crawley council aims to secure SHDF Wave 2 funding for £3m. If successful, q 0.000008866

Whilst the measures stated above and in Table 2.2 will help to contribute towards compliance, Crawley Borough Council anticipates that further additional measures not yet prescribed will be required in subsequent years to achieve compliance and enable the revocation of Crawley AQMA.

Measure No.	Measure	Category	Classification	Year Measure Introduced	Estimated / Actual Completion Year	Organisations Involved	**Funding S1 0 G( )Qr**

Measure No.	Measure	Category	Classification	Year Measure Introduced	Estimated / Actual Completion Year	Organisations Involved	**Funding Source**	**Defra AQ Grant Funding**	Funding Status	**Estimated Cost of Measure**	Measure Status	Reduction in Pollutant / Emission from Measure	K

Measure No.	Measure	Category	Classification	Year Measure Introduced	Estimated / Actual Completion Year	Organisations Involved	**Funding Source**	**Defra AQ Grant Funding**	Funding Status	**Estimated Cost of Measure**	Measure Status	Reduction in Pollutant / Emission from Measure	Key Performance Indicator	Progress to Date	Comments / Barriers to Implementation
	quality in our local area"													organise the event – but is to resume 2022.	
[redacted]	on rds for ed Taxis	Promoting Low Emissions Transport	Taxi emission incentives	2020	Due to be adopted 2021/22 and implemented from 2022/23 Delayed (see Comments)	CBC	Individual operators' capital replacement/ alternative funding sources	No	Fully / or partially grant funded	Not Known (Renewal purchase costs on individual basis)	2020 Policy update not formally adopted due to Covid impact on taxi trade. Outcome of 2021 review of policy/ consultation with taxi trade: Policy was updated 1st April 2022, with strategies to improve vehicle emissions and general standards but not currently to adopt new licensed taxis zero emission capable (ZEC) from 2022	Target still for Zero emissions by 2030	Following post Covid consultation Original indicators: new taxis zero emission capable (ZEC) from Q2 2022/ existing taxis retrofitted to Euro 6 Q4 2022 Diesel phased out 2027 dropped in favour of all new taxis to meet Euro 6 from Q2 2022	Changes to policy on emission standards agreed at licensing committee April 2022. Further work to include more robust emissions strategy as part of ongoing Policy review will be taken to Licensing Committee in Q4 2022, covering engine size, emissions and vehicle age limits	

2.3 $PM_{2.5}$ – Local Authority Approach to Reducing Emissions and/or Concentrations

As detailed in Policy Guidance LAQM.PG16 (Chapter 7), local authorities are expected to work towards reducing emissions and/or concentrations of $PM_{2.5}$ (particulate matter with an aerodynamic diameter of 2.5µm or less). There is clear evidence that $PM_{2.5}$ has a significant impact on human health, including premature mortality, allergic reactions, and cardiovascular diseases.

Those most at risk from air pollution are the young and elderly and those with predisposed medical conditions, which may be exacerbated by elevated levels of air pollution.

PHE have produced a Public Health Outcomes Framework (PHOF) which identifies an indicator for the fraction of mortality attributable to particulate air pollution in each authority in the UK. Using this framework it is possible to compare the values for Crawley to regional and national values, as well as other nearby authorities in Sussex.

In Crawley, the latest (2020) estimated fraction of mortality attributable to long-term exposure to particulate pollution was 6.2% (this is the value using a new method which has adjusted up the values across the whole of the UK.).

The mortality attributable to particulate pollution in Crawley has remained the same as the previous year (5.4% comparing values calculated using the old method) indicating there has been no worsening of particulate pollution in the area.

These indicators are calculated for all local authorities in England, and Crawley's level (6.2) places it in a similar position to other urban centres in the region such as Reigate (6.2) Brighton (6), and Worthing (6), but below the higher mortality values attributable to PM in major cities such as London (7.8%)

Crawley borough council is working towards reducing $PM_{2.5}$ in our local area through measures aimed at reducing emissions from a range of sources in the area, including transport, industrial processes and domestic burning.

The council is taking the following measures to address $PM_{2.5}$:

1. Smoke Control Areas (SCA) in Crawley: Almost all of Crawley (with the exception of Gatwick Airport

smogs of the 1950s) when Crawley was a rapidly expanding new town. The regulation of smoke emissions by means of controls on the solid fuel appliances and the type of fuel burnt within the SCA, will reduce the impact of PM2.5 emissions in the borough. However, as the popularity of wood burners has increased in recent years, complaints of smoke from wood burners has also risen, due mainly to the burning of waste and unseasoned wood. The need to educate a new generation of solid fuel owners was therefore identified, and a “clean burn” campaign was launched in 2020 through a Defra grant funded project ‘Clean Burn Sussex”. The campaign aims to raise awareness and encourage the choice of cleaner fuels to reduce particulate emissions from domestic burning. The council has updated smoke control and domestic burning guidance on its website linking to the Clean Burn campaign.

2. Regulation of Industrial Process: Control emissions of $PM_{2.5}$ from mineral processes such as concrete batching, concrete crushing and road-stone coating.
3. Air Quality Action Plan: Many of the action plan measures listed in Table 2.2 promote low emission travel alternatives (e.g., cycling, walking, electric vehicles, car sharing etc) and the Crawley Growth Programme infrastructure projects help facilitate modal change, which together work to reduce particulate emissions.
4. Policy Measures: These include procurement of low emission vehicles and tightening the emissions standards for licensed taxis
5. Local Plan Policy: Requirement to adhere to the Sussex Air Quality and Emissions Mitigation Guidance document to reduce and, where that is not possible, mitigate emissions.
6. Local Transport Plan: Traffic management measures to reduce congestion, improve traffic flow and reduce road traffic pollutant emissions (including $PM_{2.5}$)
7. Monitoring: Direct monitoring of $PM_{2.5}$

3 Air Quality Monitoring Data and Comparison with Air Quality Objectives and National Compliance

This section sets out the monitoring undertaken within 2021 by Crawley Borough Council and how it compares with the relevant air quality objectives. In addition, monitoring results are presented for a five-year period between 2017 and 2021 to allow monitoring trends to be identified and discussed.

3.1 Summary of Monitoring Undertaken

3.1.1 Automatic Monitoring Sites

Crawley Borough Council undertook automatic (continuous) monitoring at one site during 2021. Table A.1 in Appendix A shows the details of the automatic monitoring sites. NB. Local authorities do not have to report annually on the following pollutants: 1,3 butadiene, benzene, carbon monoxide and lead, unless local circumstances indicate there is a problem. The Sussex-air page presents automatic monitoring results for Crawley Borough Council, with automatic monitoring results also available through the UK-Air website.

Maps showing the location of the monitoring sites are provided in Appendix D. Further details on how the monitors are calibrated and how the data has been adjusted are included in Appendix C.

3.1.2 Non-Automatic Monitoring Sites

Crawley Borough Council undertook non- automatic (i.e. passive) monitoring of NO_2 at 51 sites during 2021. Table A.2 in Appendix A presents the details of the non-automatic sites.

Maps showing the location of the monitoring sites are provided in Appendix D. Further details on Quality Assurance/Quality Control (QA/QC) for the diffusion tubes, including bias adjustments and any other adjustments applied (e.g. annualisation and/or distance correction), are included in Appendix C.

3.2 Individual Pollutants

The air quality monitoring results presented in this section are, where relevant, adjusted for bias, annualisation (where the annual mean data capture is below 75% and greater than 25%), and distance correction. Further details on adjustments are provided in Appendix C.

3.2.1 Nitrogen Dioxide (NO_2)

Table A.3 and Table A.4 in Appendix A compare the ratified and adjusted monitored NO_2 annual mean concentrations for the past five years with the air quality objective of 40µg/m^3. Note that the concentration data presented represents the concentration at the location of the monitoring site, following the application of bias adjustment and annualisation, as required (i.e. the values are exclusive of any consideration to fall-off with distance adjustment).

For diffusion tubes, the full 2021 dataset of monthly mean values is provided in Appendix B. Note that the concentration data presented in Table B.1 includes distance corrected values, only where relevant.

Table A.5 in Appendix A compares the ratified continuous monitored NO_2 hourly mean concentrations for the past five years with the air quality objective of 200µg/m^3, not to be exceeded more than 18 times per year.

The data in Table B.1 shows that air quality in Crawley is mainly good. There is no evidence that the hourly objective for NO_2 was being exceeded at any sites across Crawley in 2021 (indicator level annual means > 60µg/m^3) and the annual mean objective of 40µg/m^3 was 2 12T 0 004B>-3<004ev3(in)-3((1)t)-33(a)-323n-meets t

Fig 3.4

2021 Roadside NO_2 in Crawley

There were no exceedances of the hourly mean objectives for NO_2 at any of the roadside monitoring sites in Crawley in 2021. Three sites (CR63, CR93 and CR101) exceeded the annual mean objective of 40µg/m^3, but after fall off with distance adjustments there was no relevant public exposure (see Appendix C Table C.4).

Fig 3.5 shows the monthly trend in roadside NO_2 at three sites in Crawley with exceedances of the annual mean objective of 40µg/m^3 in 2021.

Fig 3.5

Fig 3.8 CR93 A2220 Three Bridges



Fig 3.9 CR60 A2219 Pegler Way



Most roadside sites in the borough saw an average drop in NO_2 levels of about 20% in 2020 compared to 2019 pre-covid levels as a direct result of fewer vehicles on the roads. By 2021 NO_2 at most roadside sites was increasing, but still below pre-covid levels, with some remaining at 2020 levels.

The CR63 monitoring site at the Hazelwick roundabout on A2011 Crawley Avenue (Fig 3.7) showed a smaller NO_2 reduction of 10% in 2020, and by 2021 NO_2 had returned to

Fig 3.11 Three Bridges Station in relation to CR93 Monitoring Site, A2220



2021 NO_2 in Crawley AQMA

There were no exceedances of the annual or hourly mean objectives for NO_2 at sites with relevant exposure within the AQMA in 2021, but CR93 close to Three Bridges station was close to exceeding the air quality limit for NO2 despite traffic levels remaining 15 % lower than at pre-Covid levels. It remains to be seen if commuter traffic to Three bridges station will return to pre-Covid levels.

Although there were no exceedeances of the AQO for NO_2 in 2021, levels continue to be high (within 10% 0f the AQO). Therefore no changes to the exisiting AQMA are proposed unless a continuing trend of reduced NO_2 concentrations is maintained in future years.

3.2.2 Particulate Matter (PM_{10})

Table A.6 in Appendix A: Monitoring Results compares the ratified and adjusted monitored PM_{10} annual mean concentrations for the past five years with the air quality objective of 40µg/m^3.

Table A.7 in Appendix A compares the ratified continuous monitored PM_{10} daily mean concentrations for the past five years with the air quality objective of 50µg/m^3, not to be exceeded more than 35 times per year.

A particulate monitor has been permanently located at Crawley's automatic monitoring station (CA2)

data to be corrected for slope by dividing by 1.06. This correction has therefore been applied to $PM_{2.5}$ data reported in in Appendix A (Table A.8)

The annual average for $PM_{2.5}$ in 2021 was 8.2ug/m3 and concentrations over the previous 5 years have all been well below the national limit value of 25µg/m^3.

The site shob9H-2(e) ET Q EMC /P <</MCID 2>> BDC q 0.000008866 55.82 Tm 0 g 0 G [( )] TJ

Diffusion Tube ID	

Diffusion Tube ID	Site Name	Site Type	

Table A.3 – Annual Mean NO_2 Monitoring Results: Automatic Monitoring (µ0.2222 RG[ )]TETQq0.04q0.00m.e02.796 3tom8W* nBT/F1 12 Tf1

Table A.4 – Annual Mean NO_2 Monitoring Results: Non-Automatic Monitoring (µg/m^3)

Site ID	X OS Grid Ref (Easting)	Y OS Grid Ref (Northing)	Site Type	Valid Data Capture for Monitoring Period (%) (1)	Valid Data Capture 2021 (%) (2)	2017	2018	2019	2020	2021

Site ID	X OS Grid Ref (Easting)	Y OS Grid Ref (Northing)	Site Type	Valid Data Capture for Monitoring Period (%) [1]	Valid Data Capture 2021 (%) [2]	2017	2018	2019	2020	2021
CR 110	526928	136356	Roadside	86	50				17	18.9
CR 111	526804	136375	Roadside	100	58				22	22.9

☒ **Diffusion tube data has been bias adjusted**

☒

Figure A.1 – Trends in Annual Mean NO_2 Concentrations

Figure A1.1 Trends in Annual Mean NO_2 Concentrations: Gatwick Airport - comparison CA2, LGW3 and RG3

Figure A1.1 presents NO_2 annual mean concentrations for continuous automatic monitoring sites at Gatwick Airport, comparing Crawley's CA2 site (located east of the runway on the eastern airport boundary close to Balcombe Road residential properties) with GAL's LGW3 site (located on-airport at South Terminal runway) and RG3 (located southwest of the runway at Poles Lane in a rural area of Crawley)

A slight upturn in annual mean NO_2 was seen in 2021, but the 5-year trend continues to be downwards. Monitoring sites RG3, LGW3 and CA2 are located on a transect across the airport from southwest to northeast following the prevailing wind direction (south-westerlies) in

The sharp falloff in airport concentrations in 2020 was a direct result of the almost total shut down of the aviation industry due to Covid which also helped to demonstrate the contribution on-airport emissions of NO_2 make to annual mean concentrations in the local area. Airport NO_2 fell to the same level as residential locations around the airport

Figure A1.4 Trends in Annual Mean NO_2 Concentrations: Background NO_2



Figure A1.4 presents the 5-year trend in NO_2 annual mean concentrations for background sites in Crawley 2017 to 2021. The long-term trend shows a reduction in NO_2, reflecting the regional and national trend as policy, technological and societal measures have helped reduce background emissions. There were no exceedances of the annual mean objective at background sites in Crawley in 2021.

Prior to Covid, a flattening of the trend in background NO_2 was emerging, however, the unusually low concentrations in 2020 due to Covid have increased the downward slope. The trend will continue to be monitored and reviewed annually through the LAQM process to see how development and traffic pressures in the borough impact backgroundNO_2 levels in the post-covid recovery period.

Figure A1.5 Trends in Annual Mean NO_2 Concentrations: Roadside NO_2

Figure A1.5 presents the 5-year trend in roadside NO_2 in Crawley 2017 to 2021. There were exceedances of the annual mean objective at three roadside sites in 2021, however, after adjustment for fall off with distance, there was no relevant public exposure. The long-term trend shows a reduction in NO_2 and a downward trend.

Table A.6 – Annual Mean PM_{10} Monitoring Results ($\mu g/m^3$)

Site ID	X OS Grid Ref (Easting)	Y OS Grid Ref (Northing)	Site Type	Valid Data Capture for Monitoring

Figure A.2 – Trends in Annual Mean PM_{10} Concentrations

Figure A3.1 Trends in Annual Mean PM_{10} Concentrations at CA2 and LGW3

Figure A3.1 presents PM_{10} annual mean concentrations for continuous automatic monitoring sites at Gatwick Airport, comparing Crawley's CA2 site, located east of runway on the eastern boundary close to residential properties on Balcombe Road, with GAL's LGW3 site, located on-airport at south terminal runway. (RG3 located southwest of the runway at Poles Lane in a rural area of Crawley has no

Table A.7 – 24-Hour Mean PM_{10} Monitoring Results, Number of PM_{10} 24-Hour Means > 50µg/m^3

Site ID	X OS Grid Ref (Easting)	Y OS Grid Ref (Northing)	Site Type	Valid Data Capture for Monitoring Period (%) (1)	Valid Data Capture 2021 (%) (2)	2017	2018	2019	2020	2021
CA2	529417	141496	Other/ Industri							

Figure A.3 – Trends in Number of 24-Hour Mean PM_{10} Results > 50µg/m^3

Figure A4.1 Trends in Annual Mean PM_{10} Concentrations: Gatwick Airport - comparison CA2 and LGW3

Figure A4.1 presents the 5 year trend in the number of 24-Hour Mean PM_{10} > 50µg/m^3 at the Gatwick East site (CA2) and Gatwick on-airport site (LGW3).

The on-airport site at the end of the runway (LGW3) shows a downwards trend and a reduction in the number of exceedances.

Site CA2 further away from the runway, to the east of the airport boundary and closer to the Balcome Road, shows an upward trend in the the number of exceedances of the 24-Hour Mean PM_{10} > 50µg/m^3 objective.

A new Fidas monitor was installed at theCA2 site March 2020 to replacing the TEOM which had been in place since 2006. For consistency all data is graphed, however, the data may not be directly comparable as prior to 2020 $PM_{2.5}$ concentrations were estimated from the CA2 TEOM measurements using local ratio of $PM_{2.5}$ to PM_{10} (Technical Guidance TG (16) methodology Box7.7).

Both sites have measured concentrations well below the national target value of 25µg/m^3 over the previous 5 years. The annual average $PM_{2.5}$ in 2021 was 8ug/m^3 at CA2 and 8ug/m^3 at LGW3, however, results remain above the WHO-recommended annual mean guideline value of 5µg/m^3.

Former Car Supermarket site, Fleming Way, Manor Royal Business District: 2 No. Storage and Distribution Warehouse Units (Class B8) – awaiting planning decision

Crompton Way, Manor Royal Business District: 3 No. Storage and Distribution Warehouse Units (Class B8) – awaiting planning decision

Land at Jersey Farm, Manor Royal Business District: 1 No. Storage and Distribution Warehouse Units (Class B8) – planning permission given but construction not started

Manor Royal District Heating Network: Two energy centres: one with ground source heat pump and one with low NOx gas boiler - feasibility study stage

Residential Development - Key Housing Sites: Sites identified as key housing sites in the Local Plan Map as planned development or currently under construction.

Residential development currently under construction:

- Zurich House, East Park, Southgate (59 dwellings) –construction started 2020 still ongoing (progress slowed due to Covid)
- Steers Lane Phase 1 (185 dwellings) – construction started 2021
- 42-44 Brighton Road (20 dwellings) – construction started 2022
- Station Gateway, Belgrave House (33 dwellings) - construction started 2022
- Station Gateway, Overline House (commercial units) - construction started 2022 (residential units awaiting planning permission due to water neutrality issue)

Residential development coming forward 2022/23:

Over the last reporting year the issue of water neutrality has emerged as an issue for residential development in the borough that must be addressed through the planning process to ensure its compliance with the Habitat Regulations. This is currently delaying the progress of most major residential development in Crawley.

Longley House (120 dwellings) – Planning permission given construction due 2022/23

Breezehusrt Drive (85 dwellings) – Planning permission given construction due 2022/23

Ambulance Station site, Ifstrenu:

Station Gateway, Mokka Site, (152 dwellings) - awaiting planning permission (delayed due to water neutrality issue)

Land East of Tinsley Lane (150 dwellings) – awaiting planning permission (delayed due to water neutrality issue)

Steers Lane Phase 2 (60 dwellings) -

including a bridge across the River Mole. As a project identified with a potential to impact air quality, the development will be subject to an EIA (as it is of a type falling within Schedule 2 of Town and Country Planning (Environmental Impact Assessment) Regulations 2017). The Scoping consultation has been completed but planning delayed due to water neutrality issue.

Gatwick Northern Runway Expansion

The Gatwick Northern Runway Expansion Development is currently going through the Consent Order (DCO) process. The Proposed Development includes alterations to the existing northern runway to provide dual runway operations and enable increased capacity at the airport. The potential passenger throughput with development is predicted to be 74 million passengers per annum (mppa) by 2038. This represents a 13mppa increase above the “without development” potential of the single runway airport.

The proposals include construction works over a 15-year period, increased on-

checked by the analyst for blank levels. If the tubes are stored for more than one week, a further sample is taken and checked for any increases in blank levels. If the levels reach a pre-determined value, the batch of tubes is discarded.

Method Calibration: A full five to seven (dependant on range of concentrations being measured) point calibration is carried out monthly using NIST certified nitrite standards. The linear graph acceptance is $r^2 = 0.999$. At the start of every batch of tubes analysed, two nitrite standards are run to check the accuracy of the calibration graph, this is repeated at the end of the analysis run. Statistical graphs are maintained using the plots of the daily standard results and the acceptance criteria achieved before an analysis run is made. An

The national bias adjustment factor is 0.84 which has been obtained from the spreadsheet version 06/22, for Gradko diffusion tubes (20% TEA in water) using 34 studies.



The local bias adjustment factor was derived from the co-located diffusion tubes (prepared and analysed by Gradko) at the Gatwick East continuous analyser site (CA2). Details of the co-location study are present in below.



The 2021 local bias correction for Crawley was calculated as 0.96 using the LAQM Diffusion Tube Precision Accuracy Spreadsheet tool. A summary table showing the calculation of the local bias adjustment factor is given in Table C,3 below. Crawley Borough Council has applied a single local bias adjustment factor of 0.96 to adjust the

and local factors were relatively close in value, and the precision and accuracy of the local co-location study was good, the more cautious approach would be to use the local factor. Consequently all conclusions and recommendations made in this report were based on monitoring results adjusted with the 2021 bias adjustment figure of 0.96.

Table C.1 – Bias Adjustment Factor

Monitoring Year	Local or National	If National , Version of National Spreadsheet	Adjustment Factor
2021	Local	N/A	0.96
2020	Local	N/A	0.98
2019			

by the Management Unit (MU). The MU for Crawley's monitoring equipment is Bureau Veritas UK (Air Quality Monitoring and Management).

Verification and Ratification process,

Bureau Veritas UK carries out data verification and ratification on automatic monitoring data for the Sussex Air Quality Partnership.

Data verification and ratification is the process whereby provisional data are combined with all other relevant information to derive the best final dataset, which is as accurate as possible and has known measurement uncertainties to allow meaningful comparison with other data using specialised data handling software.

Verification is carried out on an ongoing basis and is nominally a process to "clean-up" the initial provisional data by reviewing/excluding/including any data due to instrument malfunctions or faulty calibrations, and updates to data scaling following application of the most recent calibration factors.

Ratification is a detailed manual check of the data set carried out on a monthly/ quarterly/ yearly basis. It requires a longer-term view of the dataset incorporating the results from independent QA/QC audits of the monitoring stations, and assessment on the validity of data by experienced air quality scientists. It will consider a range of variables such as: relationships between pollutants, the impact of air pollution episodes, the context of the results in the overall climate, national and regional pollutant patterns, long-term trends etc. Once all the checks and corrections have been completed the data is given a "fully ratified" status.

Service and Maintenance of the Automatic Monitoring Equipment

The NOx analyser (ML9841B) and the FIDAS 200 particulate monitor are both maintained and twice yearly serviced on contract with specialist service engineers operating BSI audited and accredited systems for quality.

The service methodology includes; pre-service inspection, visual inspection, calibration, flow check, leak check, inspection and recording of analyser internal diagnostics and communication check from remote site.

Engineer servicing or callout reports are supplied to the council as a record of the work carried out.


Diffusion Tube ID
Distance (m)
Monitoring Site to Kerb
Receptor to Kerb
NO2 Annual Mean Concentration (µg/m3)
Bias Adjusted
Background
Predicted at Receptor
Comment
CP63
7.4
37.4
42.2
22.5
31.4
Warning: your receptor is more than 20m further from the kerb than your monitor - treat result with caution

Table C.2 – Annualisation Summary (concentrations presented in µg/m^3)

Annualisation has not been required at any site in Crawley Borough Council's monitoring network during this reporting year 2021

Table C.3 – Local Bias Adjustment Calculation

Table C.4 – NO_2 Fall off With Distance Calculations (concentrations presented in µg/m^3)

Site ID	Distance (m): Monitoring Site to Kerb	Distance (m): Receptor to Kerb	Monitored Concentration (Annualised and Bias Adjusted	Background Concentration	Concentration

Appendix D: Maps of Monitoring Locations and AQMAs

Figure D.1 – Map of Crawley AQMA Boundary



Figure D.2 – Map of Non-Automatic Monitoring Sites in Crawley in relation to the AQMA

Figure D.5 Map of Diffusion sites: CR91,93 CR

Figure D.7 Map of Diffusion sites: CR1,60,66,86,87,100,108,109,110,111

Figure D.8 Map of Diffusion site: CR104

Figure D.11 Map of Diffusion sites: CR48,50,98



Figure D.12 Map of Diffusion site: CR72

Figure D.13 Map of Diffusion sites: CR105 and 106



Figure D.14 Map of Diffusion site: CR99

Figure D.15 Map of Diffusion site: CR101

Figure D.16 Map of Diffusion site: CR102

Figure D.17 Map of Diffusion site: CR80

Figure D.18 Map of Diffusion site: CR81

Appendix E: Summary of Air Quality Objectives in England

Table E.1 – Air Quality Objectives in England[7]

Pollutant	Air Quality Objective: Concentration	Air Quality Objective: Measured as
Nitrogen Dioxide (NO_2)	200µg/m^3 not to be exceeded more than 18 times a year	1-hour mean
Nitrogen Dioxide (NO_2)	40µg/m^3	Annual mean
Particulate Matter (PM_{10})	50µg/m^3, not to be exceeded more than 35 times a year	24-hour mean
Particulate Matter (PM_{10})	40µg/m^3	Annual mean
Sulphur Dioxide (SO_2)	350µg/m^3, not to be exceeded more than 24 times a year	1-hour mean
Sulphur Dioxide (SO_2)	125µg/m^3, not to be exceeded more than 3 times a year	

Glossary of Terms

Abbreviation	Description
AADT	Annual Average Daily Traffic
AQAP	Air Quality Action Plan - A detailed description of measures, outcomes, achievement dates and implementation methods, showing how the local authority intends to achieve air quality limit values'
AQMA	Air Quality Management Area – An area where air pollutant concentrations exceed / are likely to exceed the relevant air quality objectives. AQMAs are declared for specific pollutants and objectives
AQS	Air Quality Strategy
ASR	Air quality Annual Status Report
CBC	Crawley Borough Council
CGP	Crawley Growth Programme
CAZ	Clean Air Zones
Defra	Department for Environment, Food and Rural Affairs
DMRB	Design Manual for Roads and Bridges – Air quality screening tool produced by Highways England
DFT	Department for Transport
EA	Environment Agency
EFT	Emissions Factor Toolkit
EPAQS	Expert Panel on Air Quality Standards
EU	European Union
FDMS	Filter Dynamics Measurement System
FIDAS	Fine Dust A <</MCID 55>> BDC q 154.82 183.6.914 BDC q 154.82 18 Q EMC

LPTS	Local Plan Transport Strategy
NAQS	National Air Quality Strategy
NPPF	National Planning Policy Framework
NO_2	Nitrogen Dioxide
NO_x	Nitrogen Oxides
OLEV	Office for Low Emission Vehicles
PHE	Public Health England
PM_{10}	Airborne particulate matter with an aerodynamic diameter of 10µm (micrometres or microns) or less
$PM_{2.5}$	Airborne particulate matter with an aerodynamic diameter of 2.5µm or less
QA/QC	Quality Assurance and Quality Control
SAQP	Sussex Air Quality Partnership
WHO	World Health Organisation
WSCC	West Sussex County Council
ZEC	Zero Emission Capable

References

Local Air Quality Management Technical Guidance LAQM.TG22. August 2022. Published by Defra in partnership with the Scottish Government, Welsh Assembly Government and Department of the Environment Northern Ireland.
Local Air Quality Management Policy Guidance LAQM.PG22. August 2022. Published by Defra in partnership with the Scottish Government, Welsh Assembly Government and Department of the Environment Northern Ireland.

Sussex Air Quality Emissions Mitigation Guidance 2021

Draft Crawley Borough Council Local Plan 2020-2035

Crawley Growth Programme

Crawley Town Centre Regeneration Programme 2016

National bias adjustment factor spreadsheet: http://laqm.defra.gov.uk/bias-adjustment-factors/national-bias.html
Tube precision spreadsheet: www.airquality.co.uk/archive/laqm/tools/AEA_DifTPAB_v03.xls
Bureau Vitas LAQM Diffusion Tube Data Processing Tool
Public Health Outcomes Framework